Supporting guidance for Planting of Dune Grasses

This is an old version of the page

This is an old version of the page

Date published: 30 March, 2015

Date superseded: 21 December, 2015

This item helps to stabilise sand dunes. It supports transplanting dune grasses from a suitable borrow area to an eroded face to help trap blown sand.

What needs to be done?

Pull or dig up plants from a suitable borrow area within the dune system, and replant them on the eroded face. Ensure the remaining vegetation in the borrow area will continue to protect the area from erosion.

The species you can plant include marram grass (*Ammophila arenaria*), sand couchgrass (*Elymus farctus*) and lyme grass (*Leymus arenarius*). Choose species to reflect the significant species within the dune system you have chosen to manage.

Planting offset at 50 centimetres distance will give four plants per square metre.

Transplant marram grass above the mean high-water mark.

Stock, such as sheep or cattle, can cause considerable damage to the replanted site through grazing and trampling. You may need to exclude stock to encourage the dune grasses to grow and stabilise the dunes.

How to increase the success of the option?

Transplanting should not be seen as a one-off operation. Regular maintenance is required to replace unhealthy plants, apply fertilisers, prevent trampling and extend cover to adjacent areas. Work can be undertaken at any time of year, but early spring is generally considered to be the optimum to avoid frosts, storm erosion, drought or trampling. Even assuming no storm erosion, it is likely that it will take two or three years before transplants begin to thrive and spread.

Complementary options

Combine this item with the Sand-blow Fencing option to reduce wind speed across the sand surface and encourage sand deposition.

Further information

Further information is available in Scottish Natural Heritage's coastal erosion guide and the Trust for Conservation Volunteer's Sand Dunes: A Practical Handbook.